

Product datasheet

anti-Desmocollin 1 mouse monoclonal, Dsc1-U100, liquid, purified, sample

Short overview

Cat. No.	690092S
Quantity	200 µl
Concentration	50 µg/ml (10 µg)

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	Dsc1-U100
Immunogen	Synthetic peptide corresponding to sequence present within the intracellular part of human desmocollin 1 (HQTLESVKGVGQGDGTGRYATDWQC)
Formulation	PBS pH 7.4 with 0.09% sodium azide and 0.5% BSA
UniprotID	Q08554 (Human)
Synonym	Desmocollin-1, DSC1, Cadherin family member 1, Desmosomal glycoprotein 2/3, DG2/DG3, CDHF1
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	IHC, WB
Reactivity	Human, Mouse, Rat

Applications

Immunohistochemistry (IHC) - frozen	1:50-1:100 (0.5-1 µg/ml, preincubation (directly after fixation) with 0.05-0.2% Triton X-100, for 5-10 min, depending on tissue type, is recommended; without detergent pretreatment a cytoplasmic component might cross-react)
Immunohistochemistry (IHC) - paraffin	1:50-1:100 (0.5-1 µg/ml, microwave treatment recommended; a slight crossreactivity with a lymphocyte subpopulation is abolished by mild trypsinization after the microwave step)
Western Blot (WB)	Assay dependent

Background

Mouse monoclonal antibody localizes Dsc1 present in- Suprabasal layers of interfollicular epidermis- Specific cell layers (e.g. in hair follicle root sheath)- Hassall bodies (spheroidal formations of densely packed reticulum-derived cells in the thymus). Reacts


PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com

2021 November 30 / Version: 690092S/DS-041121lim | Page 1


with 111 kDa (form "a") and 103 kDa (form "b") polypeptides (desmocollin 1 splice forms, both at pI 5.2) from human epidermal desmosomes. Reactivity on cultured cell lines: positive on primary keratinocytes, however, restricted to local piles of differentiated cells. All simple epithelia are negative.

Product images




© PROGEN in cooperation with Uniklinik Heidelberg, Dr. Jochen Heß

IHC of rat skin with anti-desmocollin 1 (1:50, courtesy of J.Heß, University Hospital Heidelberg)



© PROGEN in cooperation with Uniklinik Heidelberg, Dr. Jochen Heß

anti-Desmocollin 1 mouse monoclonal, Dsc1-U100, liquid, purified, sample



© PROGEN in cooperation with Uniklinik Heidelberg, Dr. Jochen Heß

IHC of mouse skin with anti-desmocollin 1 (1:50, courtesy of J.Heß, University Hospital Heidelberg)