

# Product datasheet

# anti-Strep-tag mouse monoclonal, C23.21, lyophilized, purified

#### Short overview

 Cat. No.
 910STR

 Quantity
 25 μg

Concentration 100 µg/ml after reconstitution with 250 µl PBS

### Product description

HostMouseAntibody TypeMonoclonalIsotypeIgG1CloneC23.21

ImmunogenStrep-tag II peptide minimal sequence WSHPQFEKFormulationLyophilized; reconstitute in 250 µl sterile PBS, pH 7.4

Synomym ChR2

Conjugate Unconjugated

**Purification** Affinity chromatography

Storage before 2-8°C until indicated expiry date

reconstitution

Storage after -20°C (avoid freeze/thaw cycles)

reconstitution

Intended use Research use only

Application WB
Reactivity Strep


#### **Applications**

Western Blot (WB) 1:5,000-1:10,000 (0.02-0.01 μg/ml)

#### Background

The monoclonal C23.21 antibody recognizes Strep II. A Step-tag is commonly added to recombinant proteins and can be used for detection or purification of the tagged protein.

## **Product images**



Western blot analysis of E. coli lysate containing Strep-tagged protein with anti-Strep-tag antibody. Western blot analysis was performed on 80 ng, 40 ng, or 10 ng of E. coli lysate containing Strep-tagged protein. Cells were lysed with SDS sample buffer. The PVDF membrane was blocked with 5% dry milk in PBST for 1 h at RT. The primary antibody anti-Strep-tag mouse monoclonal, C23.21 (Cat. No. 910STRL) was diluted in blocking buffer (antibody concentration 0.02  $\mu$ g/ml) and incubated for 1 h at RT. The secondary antibody goat anti-mouse IgG polyclonal, HRP conjugate was also diluted in blocking buffer (antibody concentration 0.2  $\mu$ g/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using PierceTM ECL Western Blotting Substrate.